

C^* -ALGEBRAS ASSOCIATED WITH ALGEBRAIC CORRESPONDENCES ON THE RIEMANN SPHERE

TSUYOSHI KAJIWARA AND YASUO WATATANI

ABSTRACT. Let $p(z, w)$ be a polynomial in two variables. We call the solution of the algebraic equation $p(z, w) = 0$ the algebraic correspondence. We regard it as the graph of the multivalued function $z \mapsto w$ defined implicitly by $p(z, w) = 0$. Algebraic correspondences on the Riemann sphere $\hat{\mathbb{C}}$ give a generalization of dynamical systems of Klein groups and rational functions. We introduce C^* -algebras associated with algebraic correspondences on the Riemann sphere. We show that if an algebraic correspondence is free and expansive on a closed p -invariant subset J of $\hat{\mathbb{C}}$, then the associated C^* -algebra $\mathcal{O}_p(J)$ is simple and purely infinite.

1. INTRODUCTION

For a branched covering $\pi : M \rightarrow M$, Deaconu and Muhly [9] introduced a C^* -algebra $C^*(M, \pi)$ as the C^* -algebra of the r -discrete groupoid constructed by Renault [27]. In order to capture information of the branched points for the complex dynamical system arising from a rational function R , in [14] we introduced a slightly different C^* -algebras $\mathcal{O}_R(\hat{\mathbb{C}})$, $\mathcal{O}_R = \mathcal{O}_R(J_R)$ and $\mathcal{O}_R(F_R)$ associated with a rational function R on the Riemann sphere, the Julia set J_R and the Fatou set F_R of R . We showed that the C^* -algebras $\mathcal{O}_R(J_R)$ on the Julia set is always simple and purely infinite if the degree of R is at least two. We also studied a relation between branched points and KMS states in [12]. C. Delaroche [1] and M. Laca - J. Spielberg [20] showed that a certain boundary action of a Kleinian group on the limit set yields a simple nuclear purely infinite C^* -algebra as groupoid C^* -algebra or crossed product. Dutkay and Jorgensen study a spectral theory on Hilbert spaces built on general finite-to-one maps ([8]).

On the other hand, Sullivan discovered a dictionary between the theory of complex analytic iteration and the theory of Kleinian groups in [29]. Sullivan's dictionary shows a strong analogy between the limit set Λ_Γ of a Kleinian group Γ and the Julia set J_R of a rational function R . Therefore it is natural to generalize both Kleinian groups and rational maps. In fact there exist such objects called algebraic correspondences or holomorphic correspondences in many works by several people, for example, Bullett [4], Bullett-Penrose [5], [6] and Münzner-Rasch [24]. Let $p(z, w)$ be a polynomial in two variables. Then the solution of the algebraic equation $p(z, w) = 0$ is called the algebraic correspondence. We regard it as the graph of the multivalued function $z \mapsto w$ defined implicitly by $p(z, w) = 0$.

In this paper, we introduce C^* -algebras associated with algebraic correspondences on the Riemann sphere. We show that if an algebraic correspondence is free and expansive on a closed p -invariant subset J of $\hat{\mathbb{C}}$, then the associated C^* -algebra $\mathcal{O}_p(J)$ is simple and purely infinite. We shall show some examples and compute the K-groups of the associated C^* -algebras. For example, Let $p(z, w) = (w - z^{m_1})(w - z^{m_2}) \dots (w - z^{m_r})$ and $m_1, \dots, m_r$ are all different, where r is the number of irreducible components. Then $J := \mathbb{T}$ is a p -invariant set. Let $b = \#B(p)$ be the number of the branched points. Then we have

$$K_0(\mathcal{O}_p(\mathbb{T})) = \mathbb{Z}^b, \quad \text{and} \quad K_1(\mathcal{O}_p(\mathbb{T})) = \mathbb{Z}/(r-1)\mathbb{Z}.$$

If $m_1, m_2, \dots, m_r$ are relatively prime, then the associated C^* -algebra $\mathcal{O}_p(\mathbb{T})$ is simple and purely infinite.

Our C^* -algebras $\mathcal{O}_p(J)$ are related with C^* -algebras of irreversible dynamical systems by Exel-Vershik [10], C^* -algebras associated with subshifts by Matsumoto [21], graph C^* -algebras [19] and their generalization for topological relations by Brenken [3], topological graphs by Katsura [16], and topological quivers by Muhly and Solel [22] and by Muhly and Tomforde [23]. Some of our C^* -algebras are isomorphic to C^* -algebras associated self-similar sets [15] and Mauldin-Williams graphs [13].

2. ALGEBRAIC CORRESPONDENCES

Let $p(z, w) \in \mathbb{C}[z, w]$ be a polynomial in two variables of degree m in z and degree n in w . We shall study an algebraic function implicitly determined by the algebraic equation $p(z, w) = 0$ on the Riemann sphere $\hat{\mathbb{C}}$. Note that there exist two different ways to compactify the algebraic curve $p(z, w) = 0$. The standard construction in algebraic geometry is to consider the zeroes of a homogeneous polynomial $P(z, w, u)$ in the complex projective plane $\mathbb{CP}^2$. But we choose the second way after [5] and introduce four variables z_1, z_2, w_1, w_2 and a polynomial

$$\tilde{p}(z_1, z_2, w_1, w_2) = z_2^m w_2^n p\left(\frac{z_1}{z_2}, \frac{w_1}{w_2}\right),$$

which is separately homogeneous in z_1, z_2 and in w_1, w_2 . We identify the Riemann sphere $\hat{\mathbb{C}}$ with the complex projective line $\mathbb{CP}^1$. We denote by $[z_1, z_2]$ an element of $\mathbb{CP}^1$. Then the *algebraic correspondence* $\mathcal{C}_p$ of $p(z, w)$ on the Riemann sphere is a closed subset of $\hat{\mathbb{C}} \times \hat{\mathbb{C}}$ defined by

$$\mathcal{C}_p := \{([z_1, z_2], [w_1, w_2]) \in \hat{\mathbb{C}} \times \hat{\mathbb{C}} \mid \tilde{p}(z_1, z_2, w_1, w_2) = 0\}.$$

Then $\mathcal{C}_p$ is compact. In fact, it is a continuous image of a compact subset

$$\{(z_1, z_2, w_1, w_2) \in \mathbb{C}^4 \mid \tilde{p}(z_1, z_2, w_1, w_2) = 0 \text{ and } |z_1|^2 + |z_2|^2 + |w_1|^2 + |w_2|^2 = 1\}.$$

To simplify notation, we write it by

$$\mathcal{C}_p = \{(z, w) \in \hat{\mathbb{C}} \times \hat{\mathbb{C}} \mid p(z, w) = 0\}$$

for short if no confusion can arise. It is also convenient to consider change of variables $u = \frac{1}{z}$ or $v = \frac{1}{w}$ instead.

For example, let $R(z) = \frac{P(z)}{Q(z)}$ be a rational function with polynomials $P(z)$, $Q(z)$. Put $p(z, w) = Q(z)w - P(z)$. Then the algebraic correspondence $\mathcal{C}_p$ of $p(z, w)$ on the Riemann sphere is exactly the graph $\{(z, w) \in \hat{\mathbb{C}} \times \hat{\mathbb{C}} \mid w = R(z), z \in \hat{\mathbb{C}}\}$ of R .

Therefore we regard the algebraic correspondence $\mathcal{C}_p$ of a general polynomial $p(z, w)$ as the graph of the algebraic function $z \mapsto w$ implicitly defined by the equation $p(z, w) = 0$. Then the iteration of the algebraic function is described naturally by a sequence $z_1, z_2, z_3, \dots$ satisfying $(z_k, z_{k+1}) \in \mathcal{C}_p$ for $k = 1, 2, 3, \dots$.

Any non-zero polynomial $p(z, w) \in \mathbb{C}[z, w]$ has a unique factorization into irreducible polynomials:

$$p(z, w) = g_1(z, w)^{n_1} \cdots g_p(z, w)^{n_p}$$

where each $g_i(z, w)$ is irreducible and g_i and g_j ($i \neq j$) are prime each other.

Throughout the paper, we assume that any polynomial $p(z, w)$ we consider is reduced, that is, the above powers $n_i = 1$ for any i . We also assume that any $g_i(z, w)$ is not a function only in z or w . In particular the degree m in z and the degree n in w of $p(z, w)$ are both greater than or equal to one.

We need to recall an elementary fact as follows:

Definition (branch index). Let $p(z, w)$ be a non-zero polynomial in two variables of degree m in z and degree n in w . Then we sometimes rewrite $p(z, w)$ as

$$\begin{aligned} p(z, w) &= a_m(w)z^m + a_{m-1}(w)z^{m-1} + \cdots + a_1(w)z + a_0(w) \\ &= b_n(z)w^n + b_{n-1}(z)w^{n-1} + \cdots + b_1(z)w + b_0(z). \end{aligned}$$

Fix $w = w_0 \in \hat{\mathbb{C}}$. Then the equation $f(z) := p(z, w_0) = 0$ in $z \in \hat{\mathbb{C}}$ has m roots with multiplicities. Take any root $z = z_0$. The branch index of $p(z, w)$ at (z_0, w_0) , denoted by $e_p(z_0, w_0)$ or $e(z_0, w_0)$, is defined to be the multiplicity for the root $z = z_0$ of $f(z) = p(z, w_0) = 0$. For example, let $R(z) = \frac{P(z)}{Q(z)}$ be a rational function with polynomials $P(z)$, $Q(z)$. Put $p(z, w) = Q(z)w - P(z)$. Then the branch index $e_p(z_0, R(z_0))$ coincides with the usual branch index $e_R(z_0)$ of R at $z = z_0$.

3. ASSOCIATED C^* -ALGEBRAS

We recall Cuntz-Pimsner algebras [25]. Let A be a C^* -algebra and X be a Hilbert right A -module. We denote by $L(X)$ be the algebra of the adjointable bounded operators on X . For $\xi, \eta \in X$, the "rank one" operator $\theta_{\xi, \eta}$ is defined by $\theta_{\xi, \eta}(\zeta) = \xi(\eta|\zeta)$ for $\zeta \in X$. The closure of the linear span of rank one operators is denoted by $K(X)$. We say that X is a Hilbert bimodule (or C^* -correspondence) over A if X is a Hilbert right A -module with a homomorphism $\phi : A \rightarrow L(X)$. In this note, we assume that X is full and ϕ is injective.

Let $F(X) = \bigoplus_{n=0}^{\infty} X^{\otimes n}$ be the Fock module of X with a convention $X^{\otimes 0} = A$. For $\xi \in X$, the creation operator $T_\xi \in L(F(X))$ is defined by

$$T_\xi(a) = \xi a \quad \text{and} \quad T_\xi(\xi_1 \otimes \cdots \otimes \xi_n) = \xi \otimes \xi_1 \otimes \cdots \otimes \xi_n.$$

We define $i_{F(X)} : A \rightarrow L(F(X))$ by

$$i_{F(X)}(a)(b) = ab \quad \text{and} \quad i_{F(X)}(a)(\xi_1 \otimes \cdots \otimes \xi_n) = \phi(a)\xi_1 \otimes \cdots \otimes \xi_n$$

for $a, b \in A$. The Cuntz-Toeplitz algebra $\mathcal{T}_X$ is the C^* -subalgebra of $L(F(X))$ generated by $i_{F(X)}(a)$ with $a \in A$ and T_ξ with $\xi \in X$. Let $j_K : K(X) \rightarrow \mathcal{T}_X$ be the homomorphism defined by $j_K(\theta_{\xi, \eta}) = T_\xi T_\eta^*$. We consider the ideal $I_X := \phi^{-1}(K(X))$ of A . Let $\mathcal{J}_X$ be the ideal of $\mathcal{T}_X$ generated by $\{i_{F(X)}(a) - (j_K \circ \phi)(a); a \in I_X\}$. Then the Cuntz-Pimsner algebra $\mathcal{O}_X$ is the quotient $\mathcal{T}_X / \mathcal{J}_X$. Let $\pi : \mathcal{T}_X \rightarrow \mathcal{O}_X$ be the quotient map. Put $S_\xi = \pi(T_\xi)$ and $i(a) = \pi(i_{F(X)}(a))$. Let $i_K : K(X) \rightarrow \mathcal{O}_X$ be the homomorphism defined by $i_K(\theta_{\xi, \eta}) = S_\xi S_\eta^*$. Then $\pi((j_K \circ \phi)(a)) = (i_K \circ \phi)(a)$ for $a \in I_X$. We note that the Cuntz-Pimsner algebra $\mathcal{O}_X$ is the universal C^* -algebra generated by $i(a)$ with $a \in A$ and S_ξ with $\xi \in X$ satisfying that $i(a)S_\xi = S_{\phi(a)\xi}$, $S_\xi i(a) = S_{\xi a}$, $S_\xi^* S_\eta = i((\xi|\eta)_A)$ for $a \in A$, $\xi, \eta \in X$ and $i(a) = (i_K \circ \phi)(a)$ for $a \in I_X$. We usually identify $i(a)$ with a in A . We denote by $\mathcal{O}_X^{alg}$ the $*$ -algebra generated algebraically by A and S_ξ with $\xi \in X$. There exists an action $\gamma : \mathbb{R} \rightarrow \text{Aut } \mathcal{O}_X$ with $\gamma_t(S_\xi) = e^{it} S_\xi$, which is called the gauge action. Since we assume that $\phi : A \rightarrow L(X)$ is isometric, there is an embedding $\phi_n : L(X^{\otimes n}) \rightarrow L(X^{\otimes n+1})$ with $\phi_n(T) = T \otimes id_X$ for $T \in L(X^{\otimes n})$ with the convention $\phi_0 = \phi : A \rightarrow L(X)$. We denote by $\mathcal{F}_X$ the C^* -algebra generated by all $K(X^{\otimes n})$, $n \geq 0$ in the inductive limit algebra $\varinjlim L(X^{\otimes n})$. Let $\mathcal{F}_n$ be the C^* -subalgebra of $\mathcal{F}_X$ generated by $K(X^{\otimes k})$, $k = 0, 1, \dots, n$, with the convention $\mathcal{F}_0 = A = K(X^{\otimes 0})$. Then $\mathcal{F}_X = \varinjlim \mathcal{F}_n$.

Let $p(z, w)$ be a non-zero polynomial in two variables and $\mathcal{C}_p$ the algebraic correspondence of $p(z, w)$ on the Riemann sphere. Consider a C^* -algebra $A = C(\hat{\mathbb{C}})$ of continuous functions on $\hat{\mathbb{C}}$. Let $X = C(\mathcal{C}_p)$. Then X is a A - A bimodule by

$$(a \cdot f \cdot b)(z, w) = a(z)f(z, w)b(w)$$

for $a, b \in A$ and $f \in X$. We introduce an A -valued inner product $(| |)_A$ on X by

$$(f|g)_A(w) = \sum_{\{z \in \hat{\mathbb{C}} | (z, w) \in \mathcal{C}_p\}} e_p(z, w) \overline{f(z, w)} g(z, w)$$

for $f, g \in X$ and $w \in \hat{\mathbb{C}}$. We need branch index $e_p(z, w)$ in the formula of the inner product above. Put $\|f\|_2 = \|(f|f)_A\|_\infty^{1/2}$.

Lemma 3.1. *The above A -valued inner product is well defined, that is, $\hat{\mathbb{C}} \ni w \mapsto (f|g)_A(w) \in \mathbb{C}$ is continuous.*

Proof. If we consider $p(z, w) = a_m(w)z^m + a_{m-1}(w)z^{m-1} + \cdots + a_1(w)z + a_0(w)$, as a polynomial in z , then each coefficient $(a_k(w))_k$ is continuous in w . Then the continuity of the map $\hat{\mathbb{C}} \ni w \mapsto (f|g)_A(w) \in \mathbb{C}$ follows from the definition of the

branch index and the continuity of the roots with multiplicities of a polynomial on the Riemann sphere. See, for example, [7]. $\square$

The left multiplication of A on X gives the left action $\phi : A \rightarrow L(X)$ such that $(\phi(a)f)(z, w) = a(z)f(z, w)$ for $a \in A$ and $f \in X$.

Proposition 3.2. *Let $p(z, w)$ be a non-zero polynomial in two variables. Then $X = C(\mathcal{C}_p)$ is a full Hilbert C^* -bimodule over $A = C(\hat{\mathbb{C}})$ without completion. The left action $\phi : A \rightarrow L(X)$ is unital and faithful.*

Proof. Let m be the degree of $p(z, w)$ in z . For any $f \in X = C(\mathcal{C}_p)$, we have

$$\|f\|_\infty \leq \|f\|_2 := \left(\sup_w \sum_{\{z \in \hat{\mathbb{C}} \mid (z, w) \in \mathcal{C}_p\}} e_p(z, w) |f(z, w)|^2 \right)^{1/2} \leq \sqrt{m} \|f\|_\infty.$$

Therefore two norms $\|\cdot\|_2$ and $\|\cdot\|_\infty$ are equivalent. Since $C(\mathcal{C}_p)$ is complete with respect to $\|\cdot\|_\infty$, it is also complete with respect to $\|\cdot\|_2$.

Since $(1|1)_A(w) = \sum_{\{z \in \hat{\mathbb{C}} \mid (z, w) \in \mathcal{C}_p\}} e_p(z, w) 1 = m$, $(X|X)_A$ contains the identity I_A of A . Therefore X is full. If $a \in A$ is not zero, then there exists $x_0 \in \hat{\mathbb{C}}$ with $a(x_0) \neq 0$. Since the degree m in z of $p(z, w)$ is greater than or equal to one, there exists $w_0 \in \hat{\mathbb{C}}$ with $(x_0, w_0) \in \mathcal{C}_p$. Choose $f \in X$ with $f(x_0, w_0) \neq 0$. Then $\phi(a)f \neq 0$. Thus ϕ is faithful. $\square$

Definition. We introduce the C^* -algebra $\mathcal{O}_p(\hat{\mathbb{C}})$ associated with an algebraic correspondence $\mathcal{C}_p = \{(z, w) \in \hat{\mathbb{C}} \times \hat{\mathbb{C}} \mid p(z, w) = 0\}$ as a Cuntz-Pimsner algebra [25] of the Hilbert C^* -bimodule $X_p = C(\mathcal{C}_p)$ over $A = C(\hat{\mathbb{C}})$.

A closed subset J in $\hat{\mathbb{C}}$ is said to be p -invariant if the following conditions are satisfied: For $z, w \in \hat{\mathbb{C}}$,

- (i) $z \in J$ and $p(z, w) = 0$ implies $w \in J$,
- (ii) $w \in J$ and $p(z, w) = 0$ implies $z \in J$.

Under the condition, we can define $\mathcal{C}_p(J) = \{(z, w) \in J \times J \mid p(z, w) = 0\}$, $A = C(J)$, $X_p(J) = C(\mathcal{C}_p(J))$ similarly. Then $X_p(J)$ is a full Hilbert C^* -bimodule (C^* -correspondence) over $A = C(J)$ and the left action is unital and faithful.

We also introduce the C^* -algebra $\mathcal{O}_p(J)$ as a Cuntz-Pimsner algebra of the Hilbert C^* -bimodule $X_p(J) = C(\mathcal{C}_p(J))$ over $A = C(J)$.

We define the set $B(p)$ of "branched points" and the set $C(p)$ of "branched values".

$$B(p) := \{z \in \hat{\mathbb{C}} \mid \text{there exists } w \in \hat{\mathbb{C}} \text{ such that } p(z, w) = 0 \text{ and } e(z, w) \geq 2\}.$$

$$C(p) := \{w \in \hat{\mathbb{C}} \mid \text{there exists } z \in \hat{\mathbb{C}} \text{ such that } p(z, w) = 0 \text{ and } e(z, w) \geq 2\}.$$

In the above definitions, we may replace $e(z, w) \geq 2$ by $\frac{\partial p}{\partial z}(z, w) = 0$ after appropriate change of variables. Symmetrically we define

$$\tilde{B}(p) := \{ w \in \hat{\mathbb{C}} \mid \text{there exists } z \in \hat{\mathbb{C}} \text{ such that } p(z, w) = 0 \text{ and } \frac{\partial p}{\partial w}(z, w) = 0 \}.$$

$$\tilde{C}(p) := \{ z \in \hat{\mathbb{C}} \mid \text{there exists } w \in \hat{\mathbb{C}} \text{ such that } p(z, w) = 0 \text{ and } \frac{\partial p}{\partial w}(z, w) = 0 \}.$$

We need some known estimates of the above sets.

Lemma 3.3. *Let $p(z, w)$ be a non-zero polynomial in two variables of degree m in z and degree n in w . Then $B(p)$, $C(p)$, $\tilde{B}(p)$ and $\tilde{C}(p)$ are finite sets. More precisely we have $\#B(p) \leq 2m(m-1)n$, $\#C(p) \leq 2(m-1)n$, $\#\tilde{B}(p) \leq 2n(n-1)m$ and $\#\tilde{C}(p) \leq 2(n-1)m$.*

Proof. It follows from proposition 2 in [5] that $\#C(p) \leq 2(m-1)n$. Since $p(z, w)$ has degree m in z , we also have $\#B(p) \leq 2m(m-1)n$. The rest is symmetrically obtained. $\square$

$$\text{Let } I_X = I_{X_p(J)} = \phi^{-1}(\phi(C(J)) \cap K(X_p(J))).$$

Proposition 3.4. $I_{X_p(J)} = \{ a \in C(J) \mid a|_{B(p)} = 0 \}$

Proof. This is a direct consequence of Proposition 4.4 in [12] or [23]. $\square$

We consider Hilbert C^* -bimodules of iteration of the "algebraic function". Put $X_A^{\otimes 2} = X \otimes_A X$, $X_A^{\otimes n} = X^{\otimes n-1} \otimes_A X$.

We define the path space $\mathcal{P}_n = \mathcal{P}_n(J)$ of length n in J by

$$\mathcal{P}_n = \{ (z_1, z_2, \dots, z_{n+1}) \in J^{n+1} \mid (z_i, z_{i+1}) \in \mathcal{C}_p(J), i = 1, \dots, n \}.$$

Then $\mathcal{P}_n$ is compact, since it is a continuous image of a compact subset. We extend the branched index for paths of length n as

$$e(z_1, z_2, \dots, z_{n+1}) = e(z_1, z_2)e(z_2, z_3) \cdots e(z_n, z_{n+1}).$$

Then $C(\mathcal{P}_n)$ is a Hilbert bimodule over A by

$$(a \cdot f \cdot b)(z_1, z_2, \dots, z_{n+1}) = a(z_1)f(z_1, z_2, \dots, z_n)b(z_{n+1})$$

$$(f|g)_A(w)$$

$$= \sum_{\{(z_1, z_2, \dots, z_n) \mid (z_1, z_2, \dots, z_n, w) \in \mathcal{P}_n\}} e(z_1, z_2, \dots, z_n, w) \overline{f(z_1, z_2, \dots, z_n, w)} g(z_1, z_2, \dots, z_n, w)$$

for $a, b \in A$, $f, g \in C(\mathcal{P}_n)$.

Lemma 3.5. *The above A -valued inner product is well defined, that is, $\hat{\mathbb{C}} \ni z \mapsto (f|g)_A(w) \in \mathbb{C}$ is continuous, for any $f, g \in C(\mathcal{P}_n)$.*

Proof. It is enough to assume that $J = \hat{\mathbb{C}}$. We may and do assume that $n = 2$, because a similar argument holds for general n . We already know that $X \otimes_A X$ has an $A = C(\hat{\mathbb{C}})$ -valued inner product. Therefore for $f_1 \otimes f_2, g_1 \otimes g_2 \in X \otimes_A X$, $\hat{\mathbb{C}} \ni w \mapsto (f_1 \otimes f_2 | g_1 \otimes g_2)_A(w) \in \mathbb{C}$ is continuous. Put $f, g \in C(\mathcal{P}_2)$ by

$$f(z_1, z_2, w) = f_1(z_1, z_2)f_2(z_2, w), \quad g(z_1, z_2, w) = g_1(z_1, z_2)g_2(z_2, w).$$

Then

$$\begin{aligned} (f|g)_A(w) &= \sum_{\{(z_1, z_2) \in \mathcal{P}_1 | (z_1, z_2, w) \in \mathcal{P}_2\}} e(z_1, z_2, w) \overline{f(z_1, z_2, w)} g(z_1, z_2, w) \\ &= \sum_{\{(z_1, z_2) \in \mathcal{P}_1 | (z_1, z_2, w) \in \mathcal{P}_2\}} e(z_1, z_2) e(z_2, w) \overline{f_1(z_1, z_2) f_2(z_2, w)} g_1(z_1, z_2) g_2(z_2, w) \\ &= \sum_{\{z_2 \in \hat{\mathbb{C}} | p(z_2, w) = 0\}} e(z_2, w) \overline{f_2(z_2, w)} \left(\sum_{\{z_1 \in \hat{\mathbb{C}} | p(z_1, z_2) = 0\}} e(z_1, z_2) \overline{f_1(z_1, z_2)} g_1(z_1, z_2) \right) g_2(z_2, w) \\ &= \sum_{\{z_2 \in \hat{\mathbb{C}} | p(z_2, w) = 0\}} e(z_2, w) \overline{f_2(z_2, w)} (f_1 | g_1)_A(z_2) g_2(z_2, w) \\ &= (f_2 | (f_1 | g_1)_A g_2)_A(w) = (f_1 \otimes f_2 | g_1 \otimes g_2)_A(w). \end{aligned}$$

Hence $w \mapsto (f|g)_A(w)$ is continuous. Then for finite sums

$$f(z_1, z_2, w) = \sum_i f_{1,i}(z_1, z_2) f_{2,i}(z_2, w), \quad g(z_1, z_2, w) = \sum_i g_{1,i}(z_1, z_2) g_{2,i}(z_2, w),$$

$w \mapsto (f|g)_A(w)$ is also continuous. Put

$$C(\mathcal{P}_2)^0 = \{f \in C(\mathcal{P}_2) \mid f(z_1, z_2, w) = \sum_{\text{finite } i} f_{1,i}(z_1, z_2) f_{2,i}(z_2, w) \text{ for } f_{1,i}, f_{2,i} \in X\}.$$

Since $C(\mathcal{P}_2)^0$ is a $*$ -subalgebra of $C(\mathcal{P}_2)$ and separates two points, $C(\mathcal{P}_2)^0$ is uniformly dense in $C(\mathcal{P}_2)$. Note that uniform norm $\|\cdot\|_\infty$ and $\|\cdot\|_2$ are equivalent, because $\|\cdot\|_\infty \leq \|\cdot\|_2 \leq m^{n/2} \|\cdot\|_\infty$. For any $f, g \in C(\mathcal{P}_2)$, there exist sequences $(f_n)_n$ and $(g_n)_n$ in $C(\mathcal{P}_2)^0$ such that $f_n \rightarrow f$ and $g_n \rightarrow g$ uniformly. Since

$$\begin{aligned} & |(f|g)_A(w) - (f_n|g_n)_A(w)| \\ & \leq \sum_{\{(z_1, z_2) | (z_1, z_2, w) \in \mathcal{P}_2\}} e(z_1, z_2, w) |\overline{f(z_1, z_2, w)} g(z_1, z_2, w) - \overline{f_n(z_1, z_2, w)} g_n(z_1, z_2, w)|, \end{aligned}$$

we see that $(f_n|g_n)_A(w)$ converges to $(f|g)_A(w)$ uniformly in w . Since a uniform limit of continuous functions is continuous, $(f|g)_A(w)$ is continuous in w . $\square$

Now it is clear to check the following proposition.

Proposition 3.6. *Let $p(z, w)$ be a non-zero polynomial in two variables. Then $X = C(\mathcal{P}_n)$ is a full Hilbert bimodule over $A = C(J)$ without completion. The left action $\phi : A \rightarrow L(X)$ is unital and faithful.*

Proposition 3.7. *There exists an isometric A - A bimodule homomorphism $\varphi : X_A^{\otimes n} \rightarrow C(\mathcal{P}_n)$ such that*

$$\varphi(f_1 \otimes f_2 \otimes \cdots \otimes f_n)(z_1, z_2, \dots, z_{n+1}) = f_1(z_1, z_2) f_2(z_2, z_3) \cdots f_n(z_n, z_{n+1})$$

for $f_1, \dots, f_n \in X$.

Proof. It is easy to check φ is a well defined A - A bimodule map. The proof of Lemma 3.5 shows that φ is isometric. Since $\|\cdot\|_\infty$ and $\|\cdot\|_2$ are equivalent, φ is onto. $\square$

We need to define another compact space $\mathcal{G}_n = \mathcal{G}_n(J)$ by

$$\mathcal{G}_n = \{ (z_1, z_{n+1}) \in J^2 \mid \text{there exists } (z_1, z_2, \dots, z_{n+1}) \in \mathcal{P}_n \}.$$

Then $C(\mathcal{G}_n)$ is a Hilbert bimodule over A by

$$(a \cdot f \cdot b)(z_1, z_{n+1}) = a(z_1) f(z_1, z_n) b(z_{n+1})$$

$$(f|g)_A(w) = \sum_{\{(z_1, z_2, \dots, z_n) \mid (z_1, z_2, \dots, z_n, w) \in \mathcal{P}_n\}} e(z_1, z_2, \dots, z_n, w) \overline{f(z_1, w)} g(z_1, w)$$

for $a, b \in A$, $f, g \in C(\mathcal{G}_n)$. Define a continuous onto map $\rho : \mathcal{P}_n \rightarrow \mathcal{G}_n$ by $\rho((z_1, z_2, \dots, z_{n+1})) = (z_1, z_{n+1})$ for $(z_1, z_2, \dots, z_{n+1}) \in \mathcal{P}_n$. The it is clear that the induced map $\rho^* : C(\mathcal{G}_n) \rightarrow C(\mathcal{P}_n)$ defined by $\rho^*(f) = f \circ \rho$ is an isometric Hilbert bimodule embedding.

4. SIMPLICITY AND PURE INFINITENESS

In this section we consider a sufficient condition for a polynomial so that the associated $\mathcal{O}_p(J)$ is simple and purely infinite.

Let J be a p -invariant subset of $\hat{\mathbb{C}}$. For any subset U of J and a natural number n , we define a subset $U^{(n)}$ of J by

$$U^{(n)} = \{ w \in J \mid (z_1, z_2, \dots, z_n, w) \in \mathcal{P}_n \text{ for some } z_1 \in U, z_2 \dots z_n \in J \}.$$

Definition. Let $p(z, w)$ be a non-zero polynomial in two variables and J a p -invariant subset of $\hat{\mathbb{C}}$. Then p is said to be *expansive* on J if for any nonempty open set $U \subset J$ in J with the relative topology there exists a natural number n such that $U^{(n)} = J$.

Example. Let $R(z) = \frac{P(z)}{Q(z)}$ be a rational function with polynomials $P(z)$, $Q(z)$ and $\deg R \geq 2$. Put $p(z, w) = Q(z)w - P(z)$. Then $U^{(n)}$ is exactly $R^n(U)$. Therefore p is expansive on the Julia set J_R by [2, Theorem 4.2.5].

Example. Let $p(z, w) = z^2 + w^2 - 1$. Then $J := \{0, 1, -1\}$ is a p -invariant set and p is not expansive on J . In fact, let $U = \{0\}$, then $U^{(2n)} = \{0\}$ and $U^{(2n+1)} = \{1, -1\}$. And p is not expansive on $\hat{\mathbb{C}}$, because an open set $U := \hat{\mathbb{C}} \setminus \{0, 1, -1\}$ is p -invariant

and $U^{(n)} = U \neq \hat{\mathbb{C}}$ for any n . In general, for any polynomial, if p has a finite p -invariant set, then p is not expansive on $\hat{\mathbb{C}}$ similarly.

Example. Let $p(z, w) = z^m - w$, $m \geq 2$. Then $J := \mathbb{T}$ is a p -invariant set and p is expansive on J , because $\mathbb{T}$ is a Julia set of $w = R(z) = z^m$.

Let $p(z, w) = z - w^n$, $n \geq 2$. Then $J := \mathbb{T}$ is a p -invariant set but p is not expansive on J . In fact $U := \hat{\mathbb{C}} \setminus \{1\}$ is p -invariant and $U^{(k)} = U \neq \hat{\mathbb{C}}$ for any k .

More generally we have a criterion.

Proposition 4.1. *Let $p(z, w) = z^m - w^n$ for natural numbers m and n . Then $J := \mathbb{T}$ is a p -invariant set. And p is expansive on $\mathbb{T}$ if and only if n is not divided by m .*

Proof. Suppose that n is divided by m , so that $n = mj$ for some $j \in \mathbb{N}$. Let $U = \{z \in \mathbb{T} \mid z^m \neq 1\}$. Then for any $k \in \mathbb{N}$, 1 is not in $U^{(k)}$. In fact, if 1 were in $U^{(k)}$, then there exist $(z_1, z_2, \dots, z_k, 1) \in \mathcal{P}_k$ such that $z_1 \in U$. Hence $z_k^m = 1$ and $z_{k-1}^m = z_k^n = z_k^{mj} = 1$. We continue this argument to obtain $z_1^m = 1$. This contradicts the fact that $z_1 \in U$. Therefore p is not expansive on $\mathbb{T}$.

Nextly, suppose that n is not divided by m . Let d be the greatest common divisor of m and n . Then $m = m_0d$ and $n = n_0d$ for some natural numbers m_0 and n_0 . Since n is not divided by m , m_0 is greater than or equal to 2. We identify $\mathbb{T}$ with $\mathbb{R} \pmod{1}$ by $z = e^{2\pi i\alpha}$ and $w = e^{2\pi i\beta}$. Then $z^m - w^n = 0$ means that $m\alpha = n\beta - k$ for some integer k . Hence

$$\mathcal{C}_p(J) \cong \{([\alpha], [\beta]) \in \mathbb{R}/\mathbb{Z} \times \mathbb{R}/\mathbb{Z} \mid \beta = \frac{m}{n}\alpha + \frac{k}{n} \text{ for some integer } k\}.$$

Then $\mathcal{C}_p(J)$ has d connected components, because $m\mathbb{R} = d\mathbb{R} \pmod{n}$.

For $([\alpha], [\beta]) \in \mathcal{G}_2(J)$, there exist $k_1, k_2 \in \mathbb{Z}$ such that

$$\beta = \frac{m}{n}(\frac{m}{n}\alpha + \frac{k_1}{n}) + \frac{k_2}{n} = \frac{m^2}{n^2}\alpha + \frac{mk_1 + nk_2}{n^2}.$$

Since $m\mathbb{Z} + n\mathbb{Z} = d\mathbb{Z}$, $([\alpha], [\beta]) \in \mathcal{G}_2(J)$ if and only if there exists $k \in \mathbb{Z}$ such that $\beta = \frac{m^2}{n^2}\alpha + \frac{dk}{n^2}$. We continue in this way to get that $([\alpha], [\beta]) \in \mathcal{G}_r(J)$ if and only if there exists $k \in \mathbb{Z}$ such that $\beta = \frac{m^r}{n^r}\alpha + \frac{d^{r-1}k}{n^r}$. Since $m^r\mathbb{Z} + n^r\mathbb{Z} = d^r\mathbb{Z}$, $\mathcal{G}_r(J)$ has $\frac{d^r}{d^{r-1}} = d$ connected components. To avoid overlapping, we see only one connected component. Hence we need to cover an interval $[0, \frac{d^{r-1}}{n^r}d] = [0, \frac{d^r}{n^r}]$. Take an open interval $I = (0, \frac{1}{m_0^r} + \varepsilon)$. Since $\frac{m^r}{n^r} \frac{1}{m_0} = \frac{d^r}{n^r}$, $I^{(r)}$ contains

$$(0, \frac{d^r}{n^r}] \cup (\frac{d^r}{n^r}, \frac{2d^r}{n^r}] \cup \dots \cup (\frac{(n_0^r - 1)d^r}{n^r}, \frac{n_0^r d^r}{n^r}] \cup \{0\} = [0, 1] \pmod{\mathbb{Z}}.$$

It is also true if we replace I by a translation of I . Now for any open set $U \subset J$, there exists an open interval (a, b) with $(a, b) \subset U$. Choose a natural number r such that $|b - a| > 1/m_0^r$. Then by the preceding argument we see that $(a, b)^{(r)} = [0, 1]$. Hence $U^{(r)} = [0, 1] \pmod{\mathbb{Z}} = J$. This shows that p is expansive on $\mathbb{T}$. □

Definition. Let N be a natural number. We define the set $\text{GP}(N)$ of N -generalized periodic points by

$$\text{GP}(N) = \{w \in J \mid \exists z \in J \exists m, n \quad 0 \leq m \neq n \leq N, \exists (z, z_2, z_3, \dots, z_n, w) \in \mathcal{P}_n, \\ \exists (z, u_2, u_3, \dots, u_m, w) \in \mathcal{P}_m\}.$$

Let $R(z) = \frac{P(z)}{Q(z)}$ be a rational function with polynomials $P(z), Q(z)$. Put $p(z, w) = Q(z)w - P(z)$. Then

$$\text{GP}(N) = \cup_{n=1}^N \{w \in \hat{\mathbb{C}} \mid R^n(w) = w\}.$$

In fact, if $R^n(w) = w$ for some $n \leq N$, then it is clear that $w \in \text{GP}(N)$. Conversely let $w \in \text{GP}(N)$, then there exists z such that $w = R^n(z) = R^m(z)$ for some $0 \leq m < n \leq N$. Then $R^{n-m}(w) = w$.

Definition. A polynomial p in two variables is said to be *free* on J if for any natural number N , $\text{GP}(N)$ is a finite set.

For example, let $R(z) = \frac{P(z)}{Q(z)}$ be a rational function with polynomials $P(z), Q(z)$. Put $p(z, w) = Q(z)w - P(z)$. If $\deg R \geq 2$, then p is free on any p -invariant set J .

Lemma 4.2. *Let $p(z, w) = z^m - w^n$. Then p is free on $J = \mathbb{T}$ if and only if $m \neq n$.*

Proof. Assume that $m \neq n$. We identify $\mathbb{T}$ with $\mathbb{R}/\mathbb{Z}$ by $z = e^{2\pi i\alpha}$ and $w = e^{2\pi i\beta}$. For any natural number N , $[\beta] \in \text{GP}(N)$ if and only if there exist $[\alpha] \in \mathbb{R}/\mathbb{Z}$ and $r, s \quad 0 \leq r \neq s \leq N$ such that $([\alpha], [\beta]) \in \mathcal{G}_r(J)$ and $([\alpha], [\beta]) \in \mathcal{G}_s(J)$. Therefore there exist $k_1, k_2 \in \mathbb{Z}$ with $0 \leq k_1 \leq n^r - 1$ and $0 \leq k_2 \leq n^s - 1$ such that

$$\beta = \frac{m^r}{n^r}\alpha + \frac{d^{r-1}k_1}{n^r} = \frac{m^s}{n^s}\alpha + \frac{d^{s-1}k_2}{n^s}.$$

Since two segment with different slopes meet at most one point, $\#\text{GP}(N) \leq n^{3N}$. Hence p is free on $\mathbb{T}$.

Conversely assume that $m = n$. Then any $(z, z, \dots, z) \in J^{k+1}$ is in $\mathcal{P}_k(J)$. Hence for any natural number N , $\text{GP}(N) = \mathbb{T}$ is an infinite set. Thus p is not free on $\mathbb{T}$. $\square$

Remark. The above example is related with an example by Katsura in [17, section 4]. If m and n are relatively prime, then his example coincides with our example. If m and n are not relatively prime, then his example is different with ours. In fact our $\mathcal{P}_n(\mathbb{T})$ is not connected if m and n are not relatively prime. But they are isomorphic as bimodule.

Proposition 4.3. *Let $R_i(z) = P_i(z)/Q_i(z)$ $i = 1, \dots, r$ be rational functions with polynomials $P_i(z), Q_i(z)$. Put $p(z, w) = (Q_1(z)w - P_1(z)) \cdots (Q_r(z)w - P_r(z))$. Let $J \subset \hat{\mathbb{C}}$ be a p -invariant closed subset. Assume that each $\deg R_i \geq 2$ and $\deg R_1, \dots, \deg R_r$ are relatively prime, then p is free on J . Furthermore, if J is a Julia set for some R_i , then p is expansive on J .*

Proof. Let N be a natural number and m, n integers with $0 \leq n < m \leq N$. For $i_1, \dots, i_m, j_1, \dots, j_n = 1, 2, \dots, r$, we shall show that

$$M := \# \{z \in \hat{\mathbb{C}} \mid R_{i_m} \circ \dots \circ R_{i_1}(z) = R_{j_n} \circ \dots \circ R_{j_1}(z)\} < \infty.$$

On the contrary, assume that $M = \infty$. Then covering degrees of both sides coincide. Count the covering derees and rearrange them. Then we have

$$(\deg R_1)^{s_1} \dots (\deg R_r)^{s_r} = (\deg R_1)^{t_1} \dots (\deg R_r)^{t_r}$$

with $s_1 + \dots + s_r = m$ and $t_1 + \dots + t_r = n$. Since $\deg R_1, \dots, \deg R_r$ are relatively prime, $s_i = t_i$ for $i = 1, \dots, r$. Then $m = n$. This contradicts the fact that $n < m$. Hence $M < \infty$. Therefore $Q(m, n) := \{z \in \hat{\mathbb{C}} \mid \text{there exists } w \in \hat{\mathbb{C}} \text{ such that } (z, w) \in \mathcal{G}_m, (z, w) \in \mathcal{G}_n\}$ is a finite set. Hence

$$\text{GP}(N) = \{w \in J \mid \exists z \in J \exists m, n \quad 0 \leq n < m \leq N, \exists (z, w) \in \mathcal{G}_m, \exists (z, w) \in \mathcal{G}_n\}$$

is also a finite set. This shows that p is free on J .

It is evident that, if J is a Julia set for some R_i , then p is expansive on J . □

Example. Let m and n be natural numbers and relatively prime. Consider $p(z, w) = (w - z^m)(w - z^n)$. We note that $J = \mathbb{T}$ is the common Julia set of $w = z^m$ and $w = z^n$. Then p is free on J and expansive on J . We note that there appears a new branched point $(1, 1)$ in $\mathcal{C}_p$.

Example. Let $R_1(z) = \frac{(z^2+1)^2}{4z(z^2-1)}$ be a rational function given by Lattes. Then the Julia set $J_{R_1} = \hat{\mathbb{C}}$. Let $R_2(z) = P_2(z)/Q_2(z)$ be any rational function with odd degree. Put $p(z, w) = ((4z(z^2 - 1))w - (z^2 + 1)^2)(Q_2(z)w - P_2(z))$. Let $J = \hat{\mathbb{C}}$. Then p is expansive on J and free on J .

Example. Let $i_1, \dots, i_n, j_1, \dots, j_n$ be natural numbers. Assume that $i_k \neq 1$ or $j_k \neq 1$ for each k . Suppose that those which are not equal to 1 are relatively prime. Put $J = \mathbb{T}$. Let

$$p(z, w) = (z^{i_1} - w^{j_1})(z^{i_2} - w^{j_2}) \dots (z^{i_n} - w^{j_n}).$$

Then p is free on J .

Example. Let m be a natural number with $m \geq 2$. Put $p(z, w) = (w - z^m)(w^m - z)$. Then p is not free on $\mathbb{T}$. In fact, there exist different paths $(z, z^m, z, z^m, z) \in \mathcal{P}_4(\mathbb{T})$ and $(z, z^m, z) \in \mathcal{P}_2(\mathbb{T})$. Hence $\text{GP}(4) = \mathbb{T}$.

Example. Let $p(z, w) = z^2 + w^2 - 1$. Then p is not free on $J = \hat{\mathbb{C}}$. In fact, choose any $(z, w) \in \mathcal{C}_p$. Then there exist different paths $(z, w, z, w, z) \in \mathcal{P}_4(\hat{\mathbb{C}})$ and $(z, w, z) \in \mathcal{P}_2(\hat{\mathbb{C}})$. Hence $\text{GP}(4) = \hat{\mathbb{C}}$.

Lemma 4.4. *Suppose that p is expansive on a p -invariant subset J . Then for any non-zero positive element $a \in A$ and for any $\varepsilon > 0$ there exist $n \in \mathbb{N}$ and $f \in X^{\otimes n}$ with $(f|f)_A = I$ such that*

$$\|a\| - \varepsilon \leq S_f^* a S_f \leq \|a\|.$$

Proof. Let x_0 be a point in J with $|a(x_0)| = \|a\|$. For any $\varepsilon > 0$ there exist an open neighbourhood U of x_0 in J such that for any $x \in U$ we have $\|a\| - \varepsilon \leq a(x) \leq \|a\|$. Choose another open neighbourhood V of x_0 in J and a compact subset $K \subset J$ satisfying $V \subset K \subset U$. Since p is expansive on J , there exists $n \in \mathbb{N}$ such that $V^{(n)} = J$. We identify $X^{\otimes n}$ with $C(\mathcal{P}_n) \supset \rho^*(C(\mathcal{G}_n))$ as in the paragraph after Proposition 3.7. Define closed subsets F_1 and F_2 of $J \times J$ by

$$\begin{aligned} F_1 &= \{(z, w) \in J \times J \mid (z, w) \in \mathcal{G}_n, z \in K\}, \\ F_2 &= \{(z, w) \in J \times J \mid (z, w) \in \mathcal{G}_n, z \in U^c\}. \end{aligned}$$

Since $F_1 \cap F_2 = \emptyset$, there exists $g \in C(\mathcal{G}_n)$ such that $0 \leq g(z, w) \leq 1$ and

$$g(z, w) = \begin{cases} 1, & (z, w) \in F_1 \\ 0, & (z, w) \in F_2. \end{cases}$$

Since $V^{(n)} = J$, for any $w \in J$ there exists $z_1 \in V$ such that $(z_1, w) \in \mathcal{G}_n$. Then $(z_1, w) \in F_1$ and $g(z_1, w) = 1$. Therefore

$$\begin{aligned} (\rho^*(g)|\rho^*(g))_A(w) &= \sum_{\{(z_1, \dots, z_n) \in \mathcal{P}_{n-1} \mid (z_1, \dots, z_n, w) \in \mathcal{P}_n\}} e(z_1, \dots, z_n, w) |g(z_1, w)|^2 \\ &\geq |g(z_1, w)|^2 = 1. \end{aligned}$$

Let $b := (\rho^*(g)|\rho^*(g))_A$. Then $b(y) = (\rho^*(g)|\rho^*(g))_A(y) \geq 1$. Thus $b \in A$ is positive and invertible. We put $f := \rho^*(g)b^{-1/2} \in X^{\otimes n}$. Then

$$(f|f)_A = b^{-1/2}(g|g)_A b^{-1/2} = I.$$

For any $w \in J$ and $(z_1, w) \in \mathcal{G}_n$, if $z \in U$, then $\|a\| - \varepsilon \leq a(z)$, and if $z \in U^c$, then $f(z_1, \dots, z_n, w) = g(z, w)b^{-1/2}(w) = 0$. Therefore

$$\begin{aligned} \|a\| - \varepsilon &= (\|a\| - \varepsilon)(f|f)_A(w) \\ &= (\|a\| - \varepsilon) \sum_{\{(z_1, \dots, z_n) \in \mathcal{P}_{n-1} \mid (z_1, \dots, z_n, w) \in \mathcal{P}_n\}} e(z_1, \dots, z_n, w) |f(z_1, \dots, z_n, w)|^2 \\ &\leq \sum_{\{(z_1, \dots, z_n) \in \mathcal{P}_{n-1} \mid (z_1, \dots, z_n, w) \in \mathcal{P}_n\}} e(z_1, \dots, z_n, w) a(z_1) |f(z_1, \dots, z_n, w)|^2 \\ &= (f|af)_A(w) = S_f^* a S_f(w). \end{aligned}$$

It is clear that $S_f^* a S_f = (f|af)_A \leq \|a\|(f|f)_A = \|a\|$. □

Lemma 4.5. *Suppose that p is expansive on a p -invariant subset J . Then for any non-zero positive element $a \in A$ and for any $\varepsilon > 0$ with $0 < \varepsilon < \|a\|$, there exist $n \in \mathbb{N}$ and $u \in X^{\otimes n}$ such that*

$$\|u\|_2 \leq (\|a\| - \varepsilon)^{-1/2} \quad \text{and} \quad S_u^* a S_u = I.$$

Proof. This is exactly as same as [14, Lemma 3.5]. $\square$

A step in the proof of the main theorem is to show a certain element S in a Cuntz-Pimsner algebra is 0. It is enough to show a corresponding element T in the Toeplitz algebra is 0. Since the Toeplitz algebra acts on the Fock module and the Fock module is realized as a function space, we can calculate $Tx = 0$ concretely.

We write $A = X^{\otimes 0}$. If $a \in A$, then T_a means $\phi(a) \otimes I_n$ on $X^{\otimes n}$. The following lemma is a key of the proof of our main theorem.

Lemma 4.6. *Let i and j be integers with $i, j \geq 0$ and $i \neq j$. Take $x \in X^{\otimes i}$ and $y \in X^{\otimes j}$. Suppose that $a \in A = C(J)$ satisfies the following condition:*

For any $(z_1, z_2, \dots, z_i, w) \in \mathcal{P}_i$, $(u_1, u_2, \dots, u_j, w) \in \mathcal{P}_j$, we have $a(z_1)a(u_1) = 0$. Then we have $aT_x T_y^* a^* = 0$.

Proof. It is enough to show $T_{ax} T_{ay}^* f = 0$ for any $f \in X^{\otimes r}$, $r = 0, 1, 2, \dots$. If $r < j$, then $T_{ax} T_{ay}^* f = T_{ax} 0 = 0$. Hence we may assume that $r \geq j$ and $f = f_1 \otimes f_2$ for $f_1 \in X^{\otimes j}$, $f_2 \in X^{\otimes(r-j)}$.

$$\begin{aligned} & (T_{ax} T_{ay}^*)(f_1 \otimes f_2)(z_1, z_2, \dots, z_i, z_{i+1}, \dots, z_{i+r-j+1}) \\ &= (T_{ax}(ay|f_1)_A f_2)(z_1, z_2, \dots, z_i, z_{i+1}, \dots, z_{i+r-j+1}) \\ &= (ax \otimes (ay|f_1)_A f_2)(z_1, z_2, \dots, z_i, z_{i+1}, \dots, z_{i+r-j+1}) \\ &= a(z_1)x(z_1, \dots, z_{i+1})(ay|f_1)_A(z_{i+1})f_2(z_{i+1}, \dots, z_{i+r-j+1}) \\ &= a(z_1)x(z_1, \dots, z_{i+1}) \\ & \quad \cdot \left(\sum_{(u_1, \dots, u_j, z_{i+1}) \in \mathcal{P}_j} e(u_1, \dots, u_j, z_{i+1}) \overline{a(u_1)y(u_1, \dots, u_j, z_{i+1})} f_1(u_1, \dots, u_j, z_{i+1}) \right) \cdot \\ & \quad f_2(z_{i+1}, \dots, z_{i+r-j+1}) \\ &= a(z_1) \overline{a(u_1)} x(z_1, \dots, z_{i+1}) \\ & \quad \cdot \left(\sum_{(u_1, \dots, u_j, z_{i+1}) \in \mathcal{P}_j} e(u_1, \dots, u_j, z_{i+1}) \overline{y(u_1, \dots, u_j, z_{i+1})} f_1(u_1, \dots, u_j, z_{i+1}) \right) \cdot \\ & \quad f_2(z_{i+1}, \dots, z_{i+r-j+1}) \\ &= 0. \end{aligned}$$

$\square$

We need to prepare the following elementary fact:

Lemma 4.7. Suppose that $p(z_0, w_0) = 0$, $\frac{\partial p}{\partial z}(z_0, w_0) \neq 0$ and $\frac{\partial p}{\partial w}(z_0, w_0) \neq 0$. Then there exist an open neighbourhood U of z_0 , an open neighbourhood V of w_0 and homeomorphism $\varphi : U \rightarrow V$ such that for any $z \in U$, $w \in V$, $p(z, w) = 0$ if and only if $w = \varphi(z)$.

Lemma 4.8. Assume that p is free on J . Suppose that J has no isolated points. Let N be a natural number. Then for any non-empty open set U in J , there exist points $w_0 \in U, z_i \in J$ ($i = 1, \dots, m^N$), an open neighbourhood V of w_0 with $V \subset U$, open neighbourhoods W_i of z_i in J and homeomorphisms $\Phi_i : W_i \rightarrow V$ for $i = 1, \dots, m^N$ satisfying the following.

- (1) $W_i \cap W_j = \emptyset$ for $i \neq j$.
- (2) For any $z \in W_i$, $w \in V$, we have $(z, w) \in \mathcal{G}_N$ if and only if $w = \Phi_i(z)$, in particular $w_0 = \Phi_i(z_i)$.
- (3) For any $s \in J$ with $(z_i, s) \in \mathcal{G}_k$ for some k ($1 \leq k \leq N$), there exist an open neighbourhood $W_{i,s}$ of s and homeomorphisms $\Phi_{i,s} : W_i \rightarrow W_{i,s}$ satisfying the following : for any $z \in W_i$, $w \in W_{i,s}$, we have $(z, w) \in \mathcal{G}_k$ if and only if $w = \Phi_{i,s}(z)$.
- (4) These open neighbourhoods W_i and $W_{i,s}$ for i, s have empty intersection each other.

Proof. Let D_1 be the set of $w \in J$ satisfying that there exist $u \in J$, $z \in \text{GP}(N)$ such that $(u, w) \in \mathcal{G}_N$, $(u, z) \in \mathcal{G}_k$ for some $k = 0, 1, \dots, N$.

Since p is free on J , $\text{GP}(N)$ is a finite set. Hence D_1 is also a finite set. Consider the set D_2 of $w \in J$ satisfying that there exist $u, z \in J$ such that $(u, w) \in \mathcal{G}_N$, $(u, z) \in \mathcal{G}_k$ for some $k = 0, 1, \dots, N$ and z is in $B(p)$, $C(p)$, $\tilde{B}(p)$ or $\tilde{C}(p)$. Then D_2 is a finite set. Since $D_1 \cup D_2$ is a finite set and J has no isolated points, there exist a non-empty open set $V_0 \subset U$ such that $V_0 \subset U \setminus (D_1 \cup D_2)$. Choose $w_0 \in V_0 \subset U \setminus (D_1 \cup D_2)$. There exist distinct $z_i \in J$ for $i = 1, \dots, m^N$ such that $(z_i, w_0) \in \mathcal{G}_N$. By the Lemma 4.7, we can choose a sufficiently small non-empty open set $V \subset V_0$, non-empty open neighbourhoods W_i of z_i and homeomorphisms $\Phi_i : W_i \rightarrow V$ for $i = 1, \dots, m^N$ satisfying all the above requirements. $\square$

Proposition 4.9. Let J be a p -invariant set with no isolated points. Suppose that p is expansive and free on J . For $N \in \mathbb{N}$, for any $T \in L(X^{\otimes N})$, for any $\varepsilon > 0$, there exists $a \in A^+ = C(J)^+$ with $\|a\| = 1$ such that

$$\|\phi(a)T\|^2 \geq \|T\|^2 - \varepsilon,$$

$$aS_x S_y^* a = 0 \quad \text{for any } x \in X^{\otimes i}, \text{ for any } y \in X^{\otimes j}, 0 \leq i, j \leq N, i \neq j.$$

Proof. For $N \in \mathbb{N}$, for any $T \in L(X^{\otimes N})$, for any $\varepsilon > 0$, there exists $f \in X^{\otimes N}$ with $\|f\|_2 = 1$ such that $\|T\|^2 \geq \|Tf\|_2^2 > \|T\|^2 - \varepsilon$. Hence there exists $w_1 \in J$ such that

$$\|Tf\|_2^2 = \sum_{\{(z_1, z_2, \dots, z_N) | (z_1, z_2, \dots, z_N, w_1) \in \mathcal{P}_N\}} e(z_1, z_2, \dots, z_N, w_1) |(Tf)(z_1, z_2, \dots, z_N, w_1)|^2$$

Since the function

$$w \mapsto \sum_{\{(z_1, z_2, \dots, z_N) | (z_1, z_2, \dots, z_N, w) \in \mathcal{P}_N\}} e(z_1, z_2, \dots, z_N, w) |(Tf)(z_1, \dots, w)|^2$$

is continuous, there exists an open neighbourhood U of w_1 such that for any $w \in U$

$$\sum_{\{(z_1, z_2, \dots, z_N) | (z_1, z_2, \dots, z_N, w) \in \mathcal{P}_N\}} e(z_1, z_2, \dots, z_N, w) |(Tf)(z_1, \dots, w)|^2 > \|T\|^2 - \varepsilon$$

By Propostion 4.9, there exist points $w_0 \in U, z_i \in J$ ($i = 1, \dots, m^N$), an open neighbourhood V of w_0 with $V \subset U$, open neighbourhoods W_i of z_i in J and homeomorphisms $\Phi_i : W_i \rightarrow V$ for $i = 1, \dots, m^N$ satisfying the conditions in the lemma. Choose $b \in A = C(J)$ satisfying

$$b(w_0) = 1, \quad 0 \leq b(w) \leq 1, \quad \text{supp } b \subset V.$$

Define $a \in C(J)$ by

$$a(z) = \begin{cases} b(\Phi_i(z)) & z \in W_i \\ 0 & \text{otherwise.} \end{cases}$$

Then this function a satisfies the condition in Lemma 4.6. Therefore for any $x \in X^{\otimes i}, y \in X^{\otimes j}, 0 \leq i, j \leq N, i \neq j$ we have $aS_xS_y^*a^* = 0$.

Moreove we have

$$\begin{aligned} \|\phi(a)Tf\|_2^2 &= \sup_w \sum_{\{(z, z_2, \dots, z_N) | (z, z_2, \dots, z_N, w) \in \mathcal{P}_N\}} e(z, z_2, \dots, z_N, w) |a(z)(Tf)(z, \dots, w)|^2 \\ &\geq \sup_w \sum_{\{(z, z_2, \dots, z_N) | (z, z_2, \dots, z_N, w) \in \mathcal{P}_N\}} e(z, z_2, \dots, z_N, w) |(Tf)(z, \dots, w)b(w)|^2 \\ &\geq \sum_{\{(z, z_2, \dots, z_N) | (z, z_2, \dots, z_N, w_0) \in \mathcal{P}_N\}} e(z, z_2, \dots, z_N, w_0) |(Tf)(z, \dots, w_0)b(w_0)|^2 \\ &> \|T\|^2 - \varepsilon. \end{aligned}$$

□

It is important to recall the fact that there exists an isometric $*$ -homomorphism

$$\varphi : L(X^{\otimes N}) \supset A \otimes I^N + K(X) \otimes I^{N-1} + \dots + K(X^{\otimes N}) \rightarrow \mathcal{O}_p(J)^{\mathbf{T}}$$

as in Pimsner [25, Proposition 3.11] and Fowler-Muhly-Raeburn [11, Proposition 4.6] such that

$$\varphi(a + \theta_{x_1 \otimes \dots \otimes x_k, y_1 \otimes \dots \otimes y_k}) = a + S_{x_1} \dots S_{x_k} S_{y_k}^* \dots S_{y_1}^*.$$

To simplify notation, we put $S_x = S_{x_1} \dots S_{x_k}$ for $x = x_1 \otimes \dots \otimes x_k \in X^{\otimes k}$.

Lemma 4.10. *Let J be a p -invariant set with no isolated points. Suppose that p is expansive and free on J . Let $b = c^*c$ for some $c \in \mathcal{O}_X^{\text{alg}}$. We decompose $b = \sum_j b_j$ with $\gamma_t(b_j) = e^{ijt}b_j$. For any $\varepsilon > 0$ there exists $P \in A$ with $0 \leq P \leq I$ satisfying the following:*

- (1) $Pb_jP = 0 \quad (j \neq 0)$
- (2) $\|Pb_0P\| \geq \|b_0\| - \varepsilon$

Proof. For $x \in X^{\otimes n}$, we define $\text{length}(x) = n$ with the convention $\text{length}(a) = 0$ for $a \in A$. We write c as a finite sum $c = a + \sum_i S_{x_i} S_{y_i}^*$. Put $n = 2 \max\{\text{length}(x_i), \text{length}(y_i); i\}$. For $j > 0$, each b_j is a finite sum of terms in the form such that

$$S_x S_y^* \quad x \in X^{\otimes(k+j)}, \quad y \in X^{\otimes k} \quad 0 \leq k + j \leq n.$$

In the case when $j < 0$, b_j is a finite sum of terms in the form such that

$$S_x S_y^* \quad x \in X^{\otimes k}, \quad y \in X^{\otimes(k+|j|)} \quad 0 \leq k + |j| \leq n.$$

We shall identify b_0 with an element in $L(X^{\otimes n})$. Apply Proposition 4.9 for $T = (b_0)^{1/2}$. Then there exists $a \in A^+ = C(J)^+$ with $\|a\| = 1$ such that

$$\|\phi(a)T\|^2 \geq \|T\|^2 - \varepsilon,$$

$$aS_x S_y^* a = 0 \quad \text{for any } x \in X^{\otimes i}, \text{ for any } y \in X^{\otimes j}, 0 \leq i, j \leq N, i \neq j.$$

Define a positive operator $P = a \in A$. Then

$$\|Pb_0P\| = \|Pb_0^{1/2}\|^2 \geq \|b_0^{1/2}\|^2 - \varepsilon = \|b_0\| - \varepsilon.$$

It is evident that $Pb_jP = 0$ for $j \neq 0$. □

Since we have prepared technical lemmas addapting to our particular situation, the rest of the proof of our main theorem is a standard one.

Theorem 4.11. *Let $p(z, w)$ be a reduced non-zero polynomial in two variables with a unique factorization into irreducible polynomials: $p(z, w) = g_1(z, w) \cdots g_p(z, w)$, where each $g_i(z, w)$ is irreducible and g_i and g_j ($i \neq j$) are prime each other. We assume that any $g_i(z, w)$ is not a function only in z or w . Let J be a p -invariant set with no isolated points. Suppose that p is expansive and free on J . Then the associated C^* -algebra $\mathcal{O}_p(J)$ is simple and purely infinite.*

Proof. Let $w \in \mathcal{O}_p(J)$ be any non-zero positive element. We shall show that there exist $z_1, z_2 \in \mathcal{O}_p(J)$ such that $z_1^* w z_2 = I$. We may assume that $\|w\| = 1$. Let $E : \mathcal{O}_p(J) \rightarrow \mathcal{O}_p(J)^\alpha$ be the canonical conditional expectation onto the fixed point algebra by the gauge action α . Since E is faithful, $E(w) \neq 0$. Choose ε such that

$$0 < \varepsilon < \frac{\|E(w)\|}{4} \quad \text{and} \quad \varepsilon(\|E(w)\| - 3\varepsilon)^{-1} \leq 1.$$

There exists an element $c \in \mathcal{O}_p(J)^{alg}$ such that $\|w - c^*c\| < \varepsilon$ and $\|c\| \leq 1$. Let $b = c^*c$. Then b is decomposed as a finite sum $b = \sum_j b_j$ with $\alpha_t(b_j) = e^{ijt}b_j$. Since $\|b\| \leq 1$, $\|b_0\| = \|E(b)\| \leq 1$. By Lemma 4.10, there exists $P \in A$ with $0 \leq P \leq I$ satisfying $Pb_jP = 0$ ($j \neq 0$) and $\|Pb_0P\| \geq \|b_0\| - \varepsilon$. Then we have

$$\begin{aligned} \|Pb_0P\| &\geq \|b_0\| - \varepsilon = \|E(b)\| - \varepsilon \\ &\geq \|E(w)\| - \|E(w) - E(b)\| - \varepsilon \geq \|E(w)\| - 2\varepsilon. \end{aligned}$$

For $T := Pb_0P \in L(X^{\otimes m})$, there exists $f \in X^{\otimes m}$ with $\|f\| = 1$ such that

$$\|T^{1/2}f\|_2^2 = \|(f|Tf)_A\| \geq \|T\| - \varepsilon.$$

Hence we have $\|T^{1/2}f\|_2^2 \geq \|E(w)\| - 3\varepsilon$. Define $a = S_f^*TS_f = (f|Tf)_A \in A$. Then $\|a\| \geq \|E(w)\| - 3\varepsilon > \varepsilon$. By Lemma 4.5, there exists $n \in \mathbb{N}$ and $u \in X^{\otimes n}$ such that

$$\|u\|_2 \leq (\|a\| - \varepsilon)^{-1/2} \quad \text{and} \quad S_u^*aS_u = I.$$

Then $\|u\| \leq (\|E(w)\| - 3\varepsilon)^{-1/2}$. We have

$$\|S_f^*PwPS_f - a\| \leq \|S_f\|^2\|P\|^2\|w - b\| < \varepsilon.$$

Therefore

$$\|S_u^*S_f^*PwPS_fS_u - I\| < \|u\|^2\varepsilon \leq \varepsilon(\|E(w)\| - 3\varepsilon)^{-1} \leq 1.$$

Hence $S_u^*S_f^*PwPS_fS_u$ is invertible. Thus there exists $v \in \mathcal{O}_X$ with $S_u^*S_f^*PwPS_fS_uv = I$. Put $z_1 = S_u^*S_f^*P$ and $z_2 = PS_fS_uv$. Then $z_1wz_2 = I$. $\square$

Remark. Schweizer's theorem in [28] also implies that $\mathcal{O}_p(J)$ is simple. Our theorem gives simplicity and pure infiniteness with a direct proof.

C^* -algebra $\mathcal{O}_p(J)$ is separable and nuclear, and satisfies the Universal Coefficient Theorem. Hence the isomorphism class of C^* -algebra $\mathcal{O}_p(J)$ is completely determined by the K -group together with the class of the unit by the classification theorem by Kirchberg-Phillips [18], [26].

Example. Let m and n be natural numbers. Consider $p(z, w) = z^m - w^n$ and $J = \mathbb{T}$. If n is not divided by m , then $\mathcal{O}_p(J)$ is simple and purely infinite.

Example. Let m and n be natural numbers and relatively prime. Consider $p(z, w) = (w - z^m)(w - z^n)$ and $J = \mathbb{T}$. Then $\mathcal{O}_p(J)$ is simple and purely infinite.

Example. Let $R_1(z) = \frac{(z^2+1)^2}{4z(z^2-1)}$ be a rational function given by Lattes. Let $R_2(z) = P_2(z)/Q_2(z)$ be any rational function with odd degree. Consider

$$p(z, w) = ((4z(z^2 - 1))w - (z^2 + 1)^2)(Q_2(z)w - P_2(z)).$$

Let $J = \hat{\mathbb{C}}$. Then $\mathcal{O}_p(J)$ is simple and purely infinite.

Example. Let $i_1, \dots, i_n, j_1, \dots, j_n$ be natural numbers. Assume that $i_k \neq 1$ or $j_k \neq 1$ for each k . Suppose that those which are not equal to 1 are relatively prime. Put $J = \mathbb{T}$. Let

$$p(z, w) = (z^{i_1} - w^{j_1})(z^{i_2} - w^{j_2}) \cdots (z^{i_n} - w^{j_n}).$$

Then $\mathcal{O}_p(J)$ is simple and purely infinite.

5. K-GROUPS

We shall compute K-groups for several examples.

Example. Let $p(z, w) = z^m - w^n$. Then $J := \mathbb{T}$ is a p -invariant set. Consider the Hilbert bimodule $X_p = C(\mathcal{C}_p)$ over $A = C(\mathbb{T})$. Then X_p is isomorphic to A^m as a right A -module. In fact, let $u_i(z, w) = \frac{1}{\sqrt{m}} z^i$ for $i = 0, 1, \dots, m-1$. Then $(u_i|u_j)_A = \delta_{i,j}I$ and $\{u_0, u_1, \dots, u_{m-1}\}$ is a basis of X_p , in the sense that $f = \sum_{i=0}^{m-1} u_i(u_i|f)_A$ for any $f \in X_p = C(\mathcal{C}_p)$. Let $a_1(z) = z$ for $z \in \mathbb{T}$. Then

$$(\phi(a_1)u_i)(z, w) = \frac{1}{\sqrt{m}} z^{i+1} = u_{i+1}(z, w)$$

for $i = 0, 1, \dots, m-2$. And

$$(\phi(a_1)u_{m-1})(z, w) = \frac{1}{\sqrt{m}} z^m = \frac{1}{\sqrt{m}} w^n = (u_0 \cdot a_1^n)(z, w).$$

Therefore, if we identify $\phi(a) \in L(X_p) = M_n(A)$, then $\phi(a_1)_{i,j} = I$ for $i = j+1, j = 1, \dots, m-2$, $\phi(a_1)_{0,m-1} = a_1^n$ and $\phi(a_1)_{i,j} = 0$ for others. Let $\phi_1^* : K_1(A) = \mathbb{Z} \rightarrow K_1(A) = \mathbb{Z}$. Since $[a_1]$ is the generator of $K_1(A) = \mathbb{Z}$ and $\phi_1([a_1]) = [a_1^n]$, $\phi_1(k) = nk$ for $k \in \mathbb{Z}$.

Since $\phi(I_A) = I_{M_m(A)}$, $\phi_0^* : K_0(A) = \mathbb{Z} \rightarrow K_0(A) = \mathbb{Z}$ is given by $\phi_1^*(k) = mk$ for $k \in \mathbb{Z}$. By a six-term exact sequence due to Pimsner [25], we have

$$\begin{array}{ccccc} \mathbb{Z} & \xrightarrow{id-m\cdot} & \mathbb{Z} & \xrightarrow{i_*} & K_0(\mathcal{O}_p(\mathbb{T})) \\ \delta_1 \uparrow & & & & \downarrow \delta_0 \\ K_1(\mathcal{O}_p(\mathbb{T})) & \xleftarrow{i_*} & 0 & \xleftarrow{id-n\cdot} & \mathbb{Z} \end{array}$$

Therefore

(1) $n = 1$ and $m = 1$:

$$K_0(\mathcal{O}_p(\mathbb{T})) \cong \mathbb{Z} \oplus \mathbb{Z}, \quad K_1(\mathcal{O}_p(\mathbb{T})) \cong \mathbb{Z} \oplus \mathbb{Z}.$$

(2) $n = 1$ and $m \neq 1$:

$$K_0(\mathcal{O}_p(\mathbb{T})) \cong \mathbb{Z} \oplus \mathbb{Z}/(m-1)\mathbb{Z}, \quad K_1(\mathcal{O}_p(\mathbb{T})) \cong \mathbb{Z}.$$

(3) $n \neq 1$ and $m = 1$:

$$K_0(\mathcal{O}_p(\mathbb{T})) \cong \mathbb{Z}, \quad K_1(\mathcal{O}_p(\mathbb{T})) \cong \mathbb{Z} \oplus \mathbb{Z}/(n-1)\mathbb{Z}.$$

(4) $n \neq 1$ and $m \neq 1$:

$$K_0(\mathcal{O}_p(\mathbb{T})) \cong \mathbb{Z}/(m-1)\mathbb{Z}, \quad K_1(\mathcal{O}_p(\mathbb{T})) \cong \mathbb{Z}/(n-1)\mathbb{Z}.$$

Example. Let $p(z, w) = (w - z^m)(w - z^n)$ with $(2 \leq m < n)$. Then $J := \mathbb{T}$ is a p -invariant set. Since the set $B(p)$ of branched points in $\mathcal{C}_p$ is non-empty, we need to be careful to compute the K-groups $K_0(\mathcal{O}_p(\mathbb{T}))$ and $K_1(\mathcal{O}_p(\mathbb{T}))$.

If z is a branched point, then $w = z^m = z^n$, so that $z^{n-m} = 1$. Hence $B(p) = \{1, \alpha, \alpha^2, \dots, \alpha^{n-m-1}\}$, where $\alpha = e^{2\pi i/(n-m)}$ is a primitive $(n-m)$ -th root of unity. Put

$$D(p) = \{(z, w) \in \mathcal{C}_p \mid e(z, w) \geq 2\} = \{((1, 1), (\alpha, \alpha^m), \dots, (\alpha^{n-m-1}, \alpha^{m(n-m-1)}))\}$$

and any branch index $e(\alpha^k, \alpha^{mk}) = 2$. Let $p_1(z, w) = (w - z^m)$ and $p_2(z, w) = (w - z^n)$. Let $X = C(\mathcal{C}_p)$ and

$$Y = C(\mathcal{C}_{p_1}) \oplus C(\mathcal{C}_{p_2}) = C(\{(1, z, w) \mid p_1(z, w) = 0\} \cup \{(2, z, w) \mid p_2(z, w) = 0\}).$$

We shall embedd $X = C(\mathcal{C}_p)$ into Y as bimodule over $A = C(\mathbb{T})$ by identifying the points corrsopioing to the branched points of p . Let

$$\begin{aligned} Z &:= \{f \in Y \mid f(1, z, w) = f(2, z, w), (z, w) \in D(p)\} \\ &= \{f \in Y \mid f(1, \alpha^r, \alpha^{mr}) = f(2, \alpha^r, \alpha^{mr}), \quad r = 0, 1, \dots, n - m - 1\}. \end{aligned}$$

Then Z is a closed submodule of Y and we can identify X with Z as bimodule.

We introduce a basis $\{u_1, \dots, u_{m+n}\}$ of Y as follows:

For $i = 1, \dots, m$,

$$u_i(1, z, w) = \frac{1}{\sqrt{m}} z^{i-1}, \quad u_i(2, z, w) = 0,$$

and for $i = m + 1, \dots, m + n$,

$$u_i(1, z, w) = 0, \quad u_i(2, z, w) = \frac{1}{\sqrt{n}} z^{i-m-1}.$$

Then $(u_i | u_j)_A = \delta_{i,j} I$. Therefore we can identify $f \in Y$ with $(f_i)_i \in A^{m+n}$ by

$$f_i = (u_i | f)_A, \quad f(k, z, w) = \sum_{i=1}^{m+n} u_i(k, z, w) f_i(w), \quad k = 1, 2.$$

We claim that $f(1, \alpha^r, \alpha^{mr}) = f(2, \alpha^r, \alpha^{mr})$ if and only if

$$\sum_{i=1}^m u_i(1, \alpha^r, \alpha^{mr}) f_i(\alpha^{mr}) = \sum_{i=m+1}^{m+n} u_i(2, \alpha^r, \alpha^{mr}) f_i(\alpha^{mr})$$

if and only if

$$\sum_{i=1}^m \frac{1}{\sqrt{m}} \alpha^{r(i-1)} f_i(\alpha^{mr}) = \sum_{i=m+1}^{m+n} \frac{1}{\sqrt{n}} \alpha^{r(i-m-1)} f_i(\alpha^{mr})$$

if and only if the corrsopioing vector $(f_1(\alpha^{mr}), \dots, f_{m+n}(\alpha^{mr})) \in \mathbb{C}^{m+n}$ is orthgonal to a vector

$$n_r := \left(\frac{1}{\sqrt{m}} 1, \frac{1}{\sqrt{m}} \alpha^r, \dots, \frac{1}{\sqrt{m}} \alpha^{r(m-1)}, -\frac{1}{\sqrt{n}} 1, -\frac{1}{\sqrt{n}} \alpha^r, \dots, -\frac{1}{\sqrt{n}} \alpha^{r(n-1)} \right) \in \mathbb{C}^{m+n}.$$

Let $C := \{\alpha^{mr} \mid r = 0, 1, \dots, n - m - 1\} = \{c_1, c_2, \dots, c_v\}$ and $c_i \neq c_j$, ($i \neq j$). For $k = 1, 2, \dots, v$, put $C(k) = \{r = 0, 1, \dots, n - m - 1 \mid \alpha^{mr} = c_k\}$. If we identify $Y = A^{m+n} = C(\mathbb{T})^{m+n}$, then

$$\begin{aligned} Z &= \{f = (f_i)_i \in A^{m+n} \mid (f_i(\alpha^{mr}))_i \text{ is orthogonal to } n_r, \text{ for } r = 0, 1, \dots, n - m - 1\} \\ &= \cap_{k=1}^v \{f = (f_i)_i \in A^{m+n} \mid (f_i(c_k))_i \text{ is orthogonal to } n_r \text{ in } \mathbb{C}^{m+n} \text{ for any } r \in B(k)\}. \end{aligned}$$

We see that for fixed k , the vectors n_r ($r \in B(k)$) are linearly independent. Therefore the subspace

$$H(k) := \{x = (x_i)_i \in \mathbb{C}^{m+n} \mid x \text{ is orthogonal to } n_r, r \in B(k)\}$$

has dimension $m + n - \#C(k) \geq m + n - (n - m) = 2m \geq 2$. Let

$$L(k) := \text{span} \{T \in B(\mathbb{C}^{m+n}) \mid T = \theta_{x,y} \text{ for some } x, y \in H(k)\}.$$

Therefore we have an identification

$$\mathcal{K}(Z) = \{f \in C(\mathbb{T}, M_{m+n}(\mathbb{C})) \mid f(c_k) \in L(k), k = 1, \dots, v\}.$$

We shall show that the canonical inclusion $i : \mathcal{K}(Z) \rightarrow \mathcal{K}(Y) \cong M_{m+n}(C(\mathbb{T}))$ induces the isomorphism

$$i_* : K_r(\mathcal{K}(Z)) \cong \mathbb{Z} \rightarrow K_r(\mathcal{K}(Y)) \cong \mathbb{Z}, \quad r = 0, 1.$$

Let

$$J = \{f \in C(\mathbb{T}, M_{m+n}(\mathbb{C})) \mid f(c_k) = 0, k = 1, \dots, v\}$$

and a finite dimensional algebra $Q = \bigoplus_{k=1}^v L(k)$. Then we have an exact sequence

$$0 \rightarrow J \rightarrow \mathcal{K}(Z) \xrightarrow{\pi} Q \rightarrow 0.$$

Consider the six-term exact sequence

$$\begin{array}{ccccc} K_0(J) = 0 & \longrightarrow & K_0(\mathcal{K}(Z)) & \xrightarrow{\pi^*} & K_0(Q) = \mathbb{Z}^v \\ \delta_1 \uparrow & & & & \downarrow \delta_0 \\ K_1(Q) = 0 & \xleftarrow{\pi^*} & K_1(\mathcal{K}(Z)) & \xleftarrow{i^*} & K_1(J) = \mathbb{Z}^v \end{array}$$

For $k = 1, \dots, v$, let $q_k \in L(k)$ be a minimal projection and we put a projection $r_k = (0, \dots, 0, q_k, 0, \dots, 0) \in Q$. Let $f_k \in \mathcal{K}(Z)$ a lift of r_k defined as a "piecewise linear" map with $f_k(c_j) = \delta_{k,j}$. Since $\delta_0([r_k]) = -[e^{2\pi i f_k}]$, we obtain that

$$\delta_0(n_1, \dots, n_v) = (n_v - n_1, n_1 - n_2, n_2 - n_3, \dots, n_{v-1} - n_v).$$

Since $\text{Im} \pi^* = \text{Ker} \delta_0 = \{(n, \dots, n) \in \mathbb{Z}^r \mid n \in \mathbb{Z}\} \cong \mathbb{Z}$ and π^* is one to one, we see $\pi^* : K_0(\mathcal{K}(Z)) \cong \mathbb{Z} \rightarrow \mathbb{Z}^r$ is given by $\pi^*(n) = (n, \dots, n)$. Since $\text{Im} \delta_0 = \text{Ker} i^*$ and i^* is onto, $i^* : K_1(J) \cong \mathbb{Z}^r \rightarrow K_1(\mathcal{K}(Z)) \cong \mathbb{Z}$ is given by $i^*(n_1, \dots, n_v) = n_1 + \dots + n_v$. Let $p \in C(\mathbb{T}, M_{m+n}(\mathbb{C}))$ be a projection such that $p(t)$ is a rank one projection for any $t \in \mathbb{T}$ and $p(c_k) \in L(k)$ for $k = 1, \dots, v$. Then $[p]$ is a generator of $K_0(\mathcal{K}(Z)) \cong \mathbb{Z}$ and also a generator of $K_0(\mathcal{K}(Y)) \cong \mathbb{Z}$. Let $c_k = e^{2\pi i \theta_k}$ with $0 \leq \theta_1 \leq \dots \leq \theta_v$. Let $u \in C(\mathbb{T}, M_{m+n}(\mathbb{C}))$ be a unitary such that $u(e^{2\pi i t}) = e^{2\pi i t / \theta_1}$ for $0 \leq t \leq \theta_1$ and $u(e^{2\pi i t}) = 1$ for others. Then $[u]$ is a generator of $K_1(\mathcal{K}(Z)) \cong \mathbb{Z}$ and also a generator of $K_1(\mathcal{K}(Y))$. Therefore we conclude that $i_* : K_r(\mathcal{K}(Z)) \cong \mathbb{Z} \rightarrow K_r(\mathcal{K}(Y)) \cong \mathbb{Z}$, $r = 0, 1$ is an isomorphism.

Since $I_X = \{f \in C(\mathbb{T}) \mid f(\alpha^k) = 0 \text{ for } k = 0, 1, m - n - 1\}$, we have $K_0(I_X) = 0$ and $K_1(I_X) = \mathbb{Z}^{m-n}$.

Therefore we can identify the left action $\phi : I_X \rightarrow K(X) = K(Z)$ with $\phi_1 \oplus \phi_2 : I_X \rightarrow K(Y) = K(C(\mathcal{C}_{p_1}) \oplus C(\mathcal{C}_{p_2}))$ on the level of K-groups. Hence $\phi^* : K_1(I_X) = \mathbb{Z}^{m-n} \rightarrow K_1(A) = \mathbb{Z}$ is given by $\phi^*(x_1, \dots, x_{m-n}) = \sum_{i=1}^{m-n} 2x_i$.

By a six-term exact sequence, we have

$$\begin{array}{ccccc} K_0(I_X) = 0 & \xrightarrow{j^* - \phi^*} & K_0(A) = \mathbb{Z} & \xrightarrow{i_*} & K_0(\mathcal{O}_p(\mathbb{T})) \\ \delta_1 \uparrow & & & & \downarrow \delta_0 \\ K_1(\mathcal{O}_p(\mathbb{T})) & \xleftarrow{i_*} & K_1(A) = \mathbb{Z} & \xleftarrow{j^* - \phi^*} & K_1(I_X) = \mathbb{Z}^{m-n} \end{array}$$

The canonical inclusion map $j : I_X \rightarrow A = C(\mathbb{T})$ induces $j^* : K_1(I_X) = \mathbb{Z}^{m-n} \rightarrow K_1(A) = \mathbb{Z}$ with $j^*(x_1, \dots, x_{m-n}) = \sum_{i=1}^{m-n} x_i$. Therefore we have

$$K_0(\mathcal{O}_p(\mathbb{T})) = \mathbb{Z}^{m-n}, \quad \text{and} \quad K_1(\mathcal{O}_p(\mathbb{T})) = 0.$$

Example. Let $p(z, w) = (w - z^{m_1})(w - z^{m_2}) \dots (w - z^{m_r})$ and $m_1, \dots, m_r$ are all different, where r is the number of irreducible components. Then $J := \mathbb{T}$ is a p -invariant set. Let $b = \#B(p)$ be the number of the branched points. By a similar calculation, we have

$$K_0(\mathcal{O}_p(\mathbb{T})) = \mathbb{Z}^b, \quad \text{and} \quad K_1(\mathcal{O}_p(\mathbb{T})) = \mathbb{Z}/(r-1)\mathbb{Z}.$$

REFERENCES

- [1] C. Anatharaman-Delaroche, *Purely infinite C^* -algebras arising from dynamical systems*, Bull. Soc. Math. France **125**(1997), 199-225
- [2] A. F. Beardon, *Iteration of rational functions* GTM 132, 1991, Springer New York.
- [3] B. Brenken, *C^* -algebras associated with topological relations*, J. Ramanujan Math. Soc., **19** (2004), 35–55.
- [4] S. Bullett, *Dynamics of quadratic correspondences*, Nonlinearity **1** (1988), 27-50.
- [5] S. Bullett and C. Penrose, *A gallery of iterated correspondences*, Experimental Mathematics **3** (1994), 85-105.
- [6] S. Bullett and C. Penrose, *Regular and limit sets for holomorphic correspondences*, Fundamenta Mathematica **167** (2001), 111-171.
- [7] F.C.Cucker and A.G.Corbalean, *An alternate proof of continuity of the roots of a polynomial*, Amer. Math. Monthly, **96**(1989), 342-345
- [8] D. E. Dutkay and P. E. T. Jorgensen, *Hilbert spaces built on a similarity and on dynamical renormalization*, to appear in J. Math. Phys., **47** (2006).
- [9] V. Deaconu and M. Muhly, *C^* -algebras associated with branched coverings*, Proc. AMS. **129** (2001), 1077-1086.
- [10] R. Exel and A. Vershik, *C^* -algebras of irreducible dynamical systems*, Canad. J. Math. **58** (2006), 39-63.
- [11] N. J. Fowler, P. S. Muhly and I. Raeburn, *Representations of Cuntz-Pimsner Algebras*, Indiana Univ. Math. J. **52** (2003), 569–605.
- [12] M. Izumi, T. Kajiwara and Y. Watatani, *KMS states and branched points*, Ergod. Th. & Dynam. Sys. **27** (2007), 18887-1918.
- [13] M. Ionescu and Y. Watatani, *C^* -algebras associated with Mauldin-Williams graphs*, to appear in Can. Math. Bull.

- [14] T. Kajiwara T. and Y. Watatani, *C*-algebras associated with complex dynamical systems*, Indiana Math. J. **54** (2005), 755-778.
- [15] T. Kajiwara and Y. Watatani, *C*-algebras associated with self-similar sets*, J. Operator Theory. **56** (2006), 225-247.
- [16] T.Katsura., *On C*-algebras associated with C*-correspondences* J. Funct, Anal. **217** (2004), 366-401.
- [17] T.Katsura., *A class of C*-algebras generalizing both graph algebras and homeomorphism C*-algebras IV, pure infiniteness* J. Funct, Anal. **254** (2008), 1161-1187.
- [18] E. Kirchberg, *The classification of purely infinite C*-algebras using Kasparov's theory*, preprint.
- [19] A.Kumjian, D.Pask, I. Raeburn and J. Renault, *Graphs, groupoids and Cuntz-Krieger algebras*, J. Funct. Anal. **144** (1997), 505-541.
- [20] M. Laca and J. Spielberg, *Purely infinite C*-algebras from boundary actions of discrete groups*, J. reine angew. Math. **480** (1996), 125-139.
- [21] K. Matsumoto, *On C*-algebras associated with subshifts*, Internat. J. Math. **8** (1997), 357-374.
- [22] P. Muhly and B. Solel, *On the Morita equivalence of tensor algebras*, Proc. London Math. Soc. **81** (2000), 113-168.
- [23] P. S. Muhly and M. Tomforde M., *Topological quivers*, Internat. J. Math. **16** (2005), 693-755.
- [24] H. F. Münzner and H.-M. Rasch, *Iterated algebraic functions and functional equations*, International J. Bifurcation and Chaos, **1** (1991), 803-822.
- [25] M. Pimsner, *A class of C*-algebras generating both Cuntz-Krieger algebras and crossed product by $\mathbb{Z}$* , Free probability theory, AMS, (1997), 189-212.
- [26] N. C. Phillips, *A classification theorem for nuclear purely infinite simple C*-algebras*, Documenta Math. (2000), **5**, 49-114.
- [27] J. Renault, *A groupoid approach to C*-algebras* Lecture Notes in Mathematics, 793. Springer, Berlin, 1980.
- [28] J. Schweizer , *Dilations of C*-correspondences and the simplicity of Cuntz-Pimsner algebras*, J. Funct Anal. **180** (2001), 404-425
- [29] D. Sullivan, *Quasiconformal homeomorphisms and dynamics I. Solution of the Fatou-Julia problem on wandering domains*, Annals of Math. **122** (1985), 401-418.

(Tsuyoshi Kajiwara) DEPARTMENT OF ENVIRONMENTAL AND MATHEMATICAL SCIENCES, OKAYAMA UNIVERSITY, TSUSHIMA, 700-8530, JAPAN

(Yasuo Watatani) DEPARTMENT OF MATHEMATICAL SCIENCES, KYUSHU UNIVERSITY, HAKOZAKI, FUKUOKA, 812-8581, JAPAN